

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A61K 31/00, 31/41, 31/415, 31/44</p>	<p>A1</p>	<p>(11) International Publication Number: WO 95/21609</p> <p>(43) International Publication Date: 17 August 1995 (17.08.95)</p>		
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; vertical-align: top; padding: 5px;"> <p>(21) International Application Number: PCT/IB95/00056</p> <p>(22) International Filing Date: 26 January 1995 (26.01.95)</p> <p>(30) Priority Data: 94810071.4 8 February 1994 (08.02.94) EP (34) Countries for which the regional or international application was filed: AT et al.</p> <p>(71) Applicant (for all designated States except US): CIBA-GEIGY AG [CH/CH]; Klybeckstrasse 141, CH-4002 Basle (CH).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): HUXLEY, Alica [SK/CH]; Margarethenstrasse 75, CH-4102 Binningen (CH). MATHIS, Georg [CH/CH]; Erachfeldstrasse 45, CH-8180 Bülach (CH).</p> <p>(74) Common Representative: CIBA-GEIGY AG; Patentabteilung, Klybeckstrasse 141, CH-4002 Basle (CH).</p> </td> <td style="width: 50%; vertical-align: top; padding: 5px;"> <p>(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).</p> <p>Published With international search report.</p> </td> </tr> </table>			<p>(21) International Application Number: PCT/IB95/00056</p> <p>(22) International Filing Date: 26 January 1995 (26.01.95)</p> <p>(30) Priority Data: 94810071.4 8 February 1994 (08.02.94) EP (34) Countries for which the regional or international application was filed: AT et al.</p> <p>(71) Applicant (for all designated States except US): CIBA-GEIGY AG [CH/CH]; Klybeckstrasse 141, CH-4002 Basle (CH).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): HUXLEY, Alica [SK/CH]; Margarethenstrasse 75, CH-4102 Binningen (CH). MATHIS, Georg [CH/CH]; Erachfeldstrasse 45, CH-8180 Bülach (CH).</p> <p>(74) Common Representative: CIBA-GEIGY AG; Patentabteilung, Klybeckstrasse 141, CH-4002 Basle (CH).</p>	<p>(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).</p> <p>Published With international search report.</p>
<p>(21) International Application Number: PCT/IB95/00056</p> <p>(22) International Filing Date: 26 January 1995 (26.01.95)</p> <p>(30) Priority Data: 94810071.4 8 February 1994 (08.02.94) EP (34) Countries for which the regional or international application was filed: AT et al.</p> <p>(71) Applicant (for all designated States except US): CIBA-GEIGY AG [CH/CH]; Klybeckstrasse 141, CH-4002 Basle (CH).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): HUXLEY, Alica [SK/CH]; Margarethenstrasse 75, CH-4102 Binningen (CH). MATHIS, Georg [CH/CH]; Erachfeldstrasse 45, CH-8180 Bülach (CH).</p> <p>(74) Common Representative: CIBA-GEIGY AG; Patentabteilung, Klybeckstrasse 141, CH-4002 Basle (CH).</p>	<p>(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).</p> <p>Published With international search report.</p>			
<p>(54) Title: TREATMENT OF NORMOTENSIVE GLAUCOMA WITH ANGIOTENSIN II ANTAGONISTS</p> <p>(57) Abstract</p> <p>The present invention describes the use of an angiotensin II antagonist or of a pharmaceutically acceptable salt thereof in the preparation of a pharmaceutical composition for the treatment of normal tension glaucoma and neurodegenerative processes.</p>				

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Treatment of normotensive glaucoma with angiotensin II antagonists

The present invention describes the use of an angiotensin II antagonist or of a pharmaceutically acceptable salt thereof in the preparation of a pharmaceutical composition for the treatment of normal tension glaucoma and neurodegenerative processes.

The term glaucoma includes symptoms of the eye which are especially to be attributed to increased intraocular pressure. Frequently, an obstruction to drainage of the aqueous humour leads to an increase in the intraocular pressure. Chronically raised intraocular pressure has a harmful effect on the optic nerve and the retina, which can terminally lead to blindness. Accordingly, for the treatment of glaucoma, active ingredients are used which are able to reduce significantly the intraocular pressure. Renin inhibitors have been mentioned, for example, as suitable agents for lowering the intraocular pressure (cf. Watkins, US 4 906 613). Increased intraocular pressure can also be treated with certain β -adrenoceptor blockers.

More recently, the phenomenon of so-called normal tension glaucoma ("low tension" or "normal tension glaucoma" is used synonymously) has now been clinically established in ophthalmology [J. Flammer, Fortschr. Ophthalmol. 87, 187(1990)]. Normal tension glaucoma is characterized by an intraocular pressure which is in the normal range, i.e. is not increased, but in which the optic disc (papilla nervi optici) is pathologically excavated and the field of vision is impaired. The pathogenetic factors are especially circulatory problems in the ocular blood vessels caused by atherosclerosis, hypotension, orthostasis, functional vasospasms and neurodegenerative factors.

Accordingly, a therapy for the treatment of normal tension glaucoma therefore starts with the possible factors involved mentioned above. Actually, this means that active ingredients which are able to guarantee or to improve the circulation in the ocular blood vessels are suitable agents for the treatment of normal tension glaucoma and of neurodegenerative factors.

It has now surprisingly been found that angiotensin II antagonists, preferably those having pronounced selectivity for the AT_1 receptor (also called A II blockers in the following), in general and already preclinically established A II blockers such as, for example, valsartan or losartan in particular, extremely improve the circulation in the eye generally and the ocular blood microcirculation in the choroidal and retinal ocular tissue especially. A II

- 2 -

blockers are also suitable for treating narrowed visual field or disorders on account of neurological pathogenesis. Such A II blockers are therefore suitable agents for treating the disease of normal tension glaucoma and neurodegenerative processes.

Such A II blockers typically have a preference of $\geq 99\%$ and preferably of $\geq 99.99\%$ for the AT_1 receptor in comparison to the AT_2 receptor.

In addition to the A II blockers mentioned, there are also mixed or dual angiotensin II antagonists. Dual angiotensin II antagonists have preferences of the same order of magnitude for the AT_1 and the AT_2 receptor, the relative extent varying greatly. A dual A II blocker having a preference for the AT_1 receptor of $\geq 50\%$, preferably $\geq 90\%$ and especially preferably of $\geq 95\%$, is also suitable for treating the disease of normal tension glaucoma.

AT_2 receptor antagonists having a preference of $> 99\%$ for the AT_2 receptor do not have the abovementioned effect on the microcirculation in the eye and are therefore unsuitable for use in the treatment of normal tension glaucoma.


In the following, if angiotensin II antagonists are being discussed, both A II blockers having the mentioned selectivity for the AT_1 receptor, and dual angiotensin II antagonists having the mentioned preference for the AT_1 receptor are meant.

The present invention therefore relates to the use of an angiotensin II antagonist or of a pharmaceutically acceptable salt thereof in the preparation of a pharmaceutical composition for the treatment of normal tension glaucoma and neurodegenerative processes.

Examples of angiotensin II antagonists include, but are not restricted to the classes of compound mentioned below and preferred individual compounds selected therefrom which have been disclosed under the patent number given in each case.

EP 253 310 describes A II blockers having the following general formula (I)


- 3 -



(I)


where the variables are as defined in the publication mentioned.

A preferred compound from EP 253 310 is DuP 753 (losartan), having the formula (Ia).



(Ia)

EP 324 377 discloses imidazole derivatives as angiotensin II antagonists having the general formula (II)




(II)

where the variables are as defined in the publication mentioned.


A preferred compound from EP 324 377 is DuP 532 having the formula (IIa).

- 4 -




(IIa)

EP 392 317 describes benzimidazoles of the formula (III) as A II blockers



where the variables are as defined in the publication mentioned.


A preferred compound from EP 392 317 is BIBS 39, having the formula (IIIa).



(IIIa)


EP 400 974 describes imidazole derivates of the formula (IV) as angiotensin II antagonists

- 5 -




where the variables are as defined in the publication mentioned.

A preferred compound from EP 400 974 is L-158,809, having the formula (IVa).




EP 403 159 describes angiotensin II antagonists of the formula (V)



where the variables are defined as in the publication mentioned.


A preferred compound from EP 403 159 is SKF 108 566 having the formula (Va).

- 6 -



(Va)


EP 412 848 describes quinoline derivatives of the formula (VI) as A II blockers




(VI)

where the variables are as defined in the publication mentioned.

Preferred compounds from EP 412 848 are ICI-D-8731 having the formula (VIa) and ICI-D-6888 having the formula (VIb).




(VIa)



(VIb)

- 7 -


Other imidazole derivatives of the formula (VII) are described as A II blockers in EP 426 021



(VII)


where the variables are as defined in the publication mentioned.

A preferred compound from EP 426 021 is FR 130 739 having the formula (VIIa).



(VIIa)

EP 434 249 describes benzofuran derivatives of the formula (VIII) as A II blockers




(VIII)


where the variables are as defined in the publication mentioned.

A preferred compound from EP 434 249 is GR 117 289 having the formula (VIIIa).

- 8 -




EP 443 983 describes angiotensin II antagonists of the formula (IX)




where the variables are as defined in the publication mentioned.

A preferred compound from EP 443 983 is CGP 48933 (valsartan) having the formula (IXa).



The PCT Application WO 91/14679 describes A II blockers of the formula (X)


- 9 -



(X)


where the variables are as defined in the publication mentioned.

A preferred compound from WO 91/14679 is SR 47 436, having the formula (Xa).



(Xa)

EP 459 136 describes benzimidazoles of the formula (XI) as angiotensin II antagonists




(XI)

where the variables are as defined in the publication mentioned.


A preferred compound of EP 459 136 is TCV 116 having the formula (XIa).

- 10 -



(XIa)


The PCT Application WO 91/17148 describes triazole derivatives of the formula (XII) as angiotensin II antagonists



(XII)

where the variables are as defined in the publication mentioned.


A preferred compound from WO 91/17148 is SC 50 560 having the formula (XIIa).



(XIIa)


EP 475 206 describes angiotensin II antagonists of the general formula (XIII)

- 11 -




where the variables are as defined in the publication mentioned.

A preferred compound from EP 475 206 is A-81988, having the formula (XIIIa).




DE 4 031 635 describes benzimidazoles of the formula (XIV) as angiotensin II antagonists



where the variables are as defined in the publication mentioned.

A preferred compound from DE 4 031 635 is BIBS-222, having the formula (XIVa).




EP 514 198 describes compounds of the formula (XV), which antagonize the effect of angiotensin II

- 12 -



where the variables are as defined in the publication mentioned.

A preferred compound from EP 514 198 is GR-138 950, having the formula (XVa).



The invention further relates to the use of one of the preferred angiotensin II antagonists (Ia) to (XVa) in the preparation of a pharmaceutical composition for the treatment of normal tension glaucoma.

The angiotensin II antagonists claimed generically (including preferably generically) and specifically in the abovementioned patent applications are hereby incorporated by reference as compounds to be used according to the invention.

All the abovementioned A II blockers are suitable for the medicinal use disclosed. Highly effective A II blockers which can antagonize the action of angiotensin II to $\geq 50\%$ at a concentration of $\leq 1\ \mu\text{M}$ are particularly preferred. A II blockers which can antagonize the action of angiotensin II to $\geq 50\%$ at a concentration of $\leq 10\ \text{nM}$ are therefore furthermore especially preferred.

Methods for the determination of the AT_1 binding on the one hand, and pharmacological tests for the quantitative determination of the angiotensin II effect on the other hand are described, for example, in: J.V. Duncia et al., *Drugs of the Future* **17**, 326(1992); or V.J. Dzau et al., *The Heart and Cardiovascular System*, pages 1631 - 1662 (1986), Raven

Press, New York, H.A. Fozzard, E. Haber, R.B. Jennings, A.M. Katz, H.E. Morgan (Editors).

The angiotensin II antagonists can be present as salts, in particular pharmaceutically acceptable salts. The angiotensin II antagonists, which have e.g. at least one basic centre, can form acid addition salts. These are formed, for example, with strong inorganic acids, such as mineral acids, e.g. sulfuric acid, a phosphoric acid or a hydrohalic acid, with strong organic carboxylic acids, such as C₁-C₄alkanecarboxylic acids which are unsubstituted or substituted e.g. by halogen, e.g. acetic acid, such as saturated or unsaturated dicarboxylic acids, e.g. oxalic, malonic, succinic, maleic, fumaric, phthalic or terephthalic acid, such as hydroxycarboxylic acids, e.g. ascorbic, glycolic, lactic, malic, tartaric or citric acid, such as amino acids, e.g. aspartic or glutamic acid, or such as benzoic acid, or with organic sulfonic acids, such as C₁-C₄alkane- or arylsulfonic acids which are unsubstituted or substituted e.g. by halogen, e.g. methane- or p-toluenesulfonic acid. Corresponding acid addition salts can also be formed which may additionally have a basic centre.

Angiotensin II antagonists which carry at least one acidic group can further form salts with bases. Suitable salts with bases are, for example, metal salts, such as alkali metal or alkaline earth metal salts, e.g. sodium, potassium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, thiomorpholine, piperidine, pyrrolidine, a mono-, di- or tri-lower alkylamine, e.g. ethyl-, tert-butyl-, diethyl-, diisopropyl-, triethyl-, tributyl- or dimethylpropylamine, or a mono-, di- or trihydroxy-lower alkylamine, e.g. mono-, di- or triethanolamine. Corresponding internal salts can furthermore be formed.

As a result of the close relationship between an angiotensin II antagonist in free form and in the form of a salt, hereinbefore and hereinafter a free compound is correspondingly and expediently to be understood, where appropriate, as also meaning a corresponding salt, or a salt is also to be understood as meaning a corresponding free compound.

The present application further relates to a pharmaceutical composition for the treatment of normal tension glaucoma, comprising a therapeutically effective amount of an angiotensin II antagonist or of a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable formulating agent which is suitable for topical application, in particular ophthalmic and systemic administration.

- 14 -

The present invention furthermore relates to an ophthalmic composition for the treatment of normal tension glaucoma, comprising a therapeutically effective amount of an angiotensin II antagonist or a pharmaceutically acceptable salt thereof.

The present invention furthermore relates to an ophthalmic composition for the treatment of normal tension glaucoma, comprising a therapeutically effective amount of (S)-N-(1-carboxy-2-methylprop-1-yl)-N-pentanoyl-N-[2'-(1H-tetrazol-5-yl)-biphenyl-4-ylmethyl]amine or of a pharmaceutically acceptable salt thereof.

An appropriate ophthalmic composition is applied to the eye topically, in particular in the form of a solution, an ointment, a gel or a solid insert. Compositions of this type contain the active compound, for example, in a range from about 0.000001 to about 5.0 % by weight, preferably from about 0.001 to about 1.0 % by weight, or in the range from about 0.01 to about 0.5 % by weight. The dose of the active compound can depend on various factors, such as administration procedure, need, age and/or individual condition.

Customary pharmaceutically acceptable adjuncts and additives, for example of the type mentioned below, known to the person skilled in the art are used for an appropriate ophthalmic composition, in particular carriers, solubilizers, tonicity-increasing agents, buffer substances, preservatives, thickeners, complexing agents and other adjuncts. Examples of additives and adjuncts of this type are found in U.S. Patent No. 5 134 124 and U.S. 4 906 613. The manufacture of such compositions is carried out in a manner known per se, i.e. the active ingredient is mixed with the appropriate adjuncts and/or additives to give corresponding ophthalmic compositions. Preferably, the active ingredient is administered in the form of eye drops, the active ingredient being dissolved, for example, in a carrier by means of a solubilizer. If desired, adjustment to the desired pH and/or buffering is carried out and if desired a tonicity-increasing agent is added. If desired, preservatives and/or other adjuncts complete an ophthalmic composition.

In order to prepare suitable formulations, the active ingredient of an angiotensin II antagonist is mixed with a carrier suitable for topical or general administration. Suitable carriers are especially water, mixtures of water and water-miscible solvents such as lower alkanols, vegetable oils or mineral oils comprising 0.5 - 5 % by weight of hydroxyethylcellulose, ethyl oleate, carboxymethylcellulose, polyvinylpyrrolidone and other non-toxic water-soluble polymers intended for ophthalmic use such as e.g. cellulose derivatives such as methylcellulose, alkali metal salts of carboxymethylcellulose, hydroxymethylcellulose, hydroxyethylcellulose and hydroxypropylcellulose, acrylates or methacrylates such as

- 15 -

salts of polyacrylic acid or ethyl acrylate, polyacrylamides, natural products such as gelatin, alginates, pectins, tragacanth, karaya gum, xanthan gum, carrageenan, agar and acacia, starch derivatives such as starch acetate and hydroxypropyl starch, and also other synthetic products such as polyvinyl alcohol, polyvinylpyrrolidone, polyvinyl methyl ether, polyethylene oxide, preferably crosslinked polyacrylic acid such as neutral carbopol or mixtures of these polymers. The concentration of the carrier is, for example, 1- to 100,000-times the active ingredient concentration.

Solvents used for the angiotensin II antagonists used are, for example, fatty acid glycerol polyglycol esters, fatty acid polyglycol esters, polyethylene glycols, glycerol ethers or mixtures of these compounds. A specific example of a particularly preferred solubilizer is a reaction product of castor oil and ethylene oxide, for example the commercial product Cremophor EL[®]. Reaction products of castor oil and ethylene oxide have proven particularly good solubilizers having an excellent ophthalmic tolerability. The concentration used primarily depends on the concentration of the active ingredient. At least sufficient is to be added that the active ingredient is brought into solution. As an example, the concentration of the solubilizer is 1- to 1000-times the active ingredient concentration.

Examples of buffer substances are acetate, ascorbate, borate, bicarbonate/carbonate, citrate, gluconate, lactate, phosphate, propionate and tris buffers. The amount of buffer substance is added, for example, which is necessary to guarantee and to maintain a physiologically tolerable pH range.

Tonicity-enhancing agents are, for example, ionic compounds, such as alkali metal or alkaline earth metal halides such as e.g. CaCl_2 , KBr, KCl, LiCl, NaI, NaBr or NaCl, or boric acid. Non-ionic tonicity-enhancing agents are e.g. urea, glycerol, sorbitol, mannitol, propylene glycol or dextrose. Sufficient tonicity-enhancing agent is added, for example, that the ready-to-use ophthalmic composition has an osmolality of about 50 to 400 mOsmol.

Examples of preservatives are quaternary ammonium salts such as cetrimide, benzalkonium chloride or benzoxonium chloride, alkylmercury salts of thiosalicylic acid such as e.g. thiomersal, phenylmercury nitrate, phenylmercury acetate or phenylmercury borate, parabens such as e.g. methylparaben or propylparaben, alcohols such as e.g. chlorobutanol, benzyl alcohol or phenylethanol, guanidine derivatives such as e.g. chlorhexidine or polyhexamethylenebiguanide, or sorbic acid. If desired, the amount of

- 16 -

preservative which is necessary to guarantee sterility is added to the ophthalmic composition.

The ophthalmic compositions can further contain non-toxic adjuncts such as, for example, emulsifiers, wetting agents or fillers such as e.g. the polyethylene glycols designated 200, 300, 400 and 600, or Carbowax designated 1000, 1500, 4000, 6000 and 10,000. Other adjuncts used if desired are mentioned below, but should not restrict the range of the possible adjuncts in any manner. In particular, they are complexing agents such as disodium EDTA or EDTA, antioxidants such as ascorbic acid, acetylcysteine, cysteine, sodium bisulfite, butylhydroxyanisole, butylhydroxytoluene or α -tocopherol acetate, stabilizers such as β -cyclodextrin, hydroxypropyl- β -cyclodextrin, thiourea, thiosorbitol, sodium dioctylsulfosuccinate or monothioglycerol, or other auxiliaries such as e.g. lauric acid sorbitol ester, triethanolamine oleate or palmitic acid esters. The amount and the nature of the adjunct added depends on the specific requirements, and as a rule is in the range from about 0.0001 to about 90 % by weight.

The present application further relates to an ophthalmic composition for the treatment of normal tension glaucoma, comprising a therapeutically effective amount of an angiotensin II antagonist or of a pharmaceutically acceptable salt thereof and another, therapeutically effective, pharmaceutical. This can be, for example, an antibiotic, β -blocker, calcium antagonist, anaesthetic, an antiinflammatory, a composition suitable for the treatment of intraocular pressure or another pharmacon.

The invention likewise relates to the use of an angiotensin II antagonist, or of a pharmaceutically acceptable salt thereof, in the preparation of a systemically administrable pharmaceutical composition for the treatment of normal tension glaucoma.

These pharmaceutical compositions are those for enteral administration, such as oral administration, and also rectal or parenteral administration to warm-blooded animals, the pharmacological active ingredient being contained on its own or together with customary pharmaceutical adjuncts. The pharmaceutical compositions contain e.g. about 0.1 % to 100 %, preferably from about 1 % to about 60 %, of the active ingredient. Pharmaceutical compositions for enteral or parenteral and also for ocular administration are e.g. those in unit dose form, such as sugar-coated tablets, tablets, capsules or suppositories, and also ampoules. These are prepared in a manner known per se, e.g. by means of conventional mixing, granulating, sugar coating, dissolving or lyophilizing processes. Pharmaceutical compositions for oral administration can thus be obtained by combining the active

ingredient with solid carriers, if desired granulating a mixture obtained, and processing the mixture or granules, if desired or necessary after addition of suitable adjuncts, to give tablets or sugar-coated tablet cores.

Suitable carriers are, in particular, fillers, such as sugars, e.g. lactose, sucrose, mannitol or sorbitol, cellulose compositions and/or calcium phosphates, e.g. tricalcium phosphate or calcium hydrogen phosphate, and also binders, such as starch pastes, using e.g. maize, wheat, rice or potato starch, gelatin, tragacanth gum, methylcellulose and/or polyvinylpyrrolidone, and, if desired, disintegrants, such as the abovementioned starches, and also carboxymethyl starches, crosslinked polyvinylpyrrolidone, agar or alginic acid or a salt thereof, such as sodium alginate. Adjuncts are primarily flow regulators and lubricants, e.g. silicic acid, talc, stearic acid or salts thereof, such as magnesium stearate or calcium stearate, and/or polyethylene glycol. Sugar-coated tablet cores are provided with suitable coatings which are, if desired, enteric-resistant, concentrated sugar solutions which if desired contain gum arabic, talc, polyvinylpyrrolidone, polyethylene glycol and/or titanium dioxide, lacquer solutions in suitable organic solvents or solvent mixtures or, for the composition of enteric-resistant coatings, solutions of suitable cellulose compositions, such as acetylcellulose phthalate or hydroxypropylmethylcellulose phthalate, being used inter alia. Colourants or pigments, e.g. for the identification or for the characterization of various doses of active ingredient, are added to the tablets or the sugar-coated tablet coatings.

Other orally administrable pharmaceutical compositions are hard capsules made of gelatin and soft, closed capsules made of gelatin and a softener, such as glycerol or sorbitol. The hard capsules can contain the active ingredient in the form of granules, e.g. in a mixture with fillers, such as lactose, binders, such as starches, and/or lubricants, such as talc or magnesium stearate, and, if desired, stabilizers. In soft capsules, the active ingredient is preferably dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin or liquid polyethylene glycols, it also being possible to add stabilizers.

Suitable rectally administrable pharmaceutical compositions are e.g. suppositories, which consist of a combination of the active compound with a suppository base. Suitable suppository bases are e.g. natural or synthetic triglycerides, paraffin hydrocarbons, polyethylene glycols and higher alkanols. Gelatin rectal capsules which contain a combination of the active compound with a base substance can also be used. Possible base substances are e.g. liquid triglycerides, polyethylene glycols and paraffin hydrocarbons.

Aqueous solutions of an active ingredient in water-soluble form, e.g. of a water-soluble salt, also suspensions of the active ingredient, such as appropriate oily injection suspensions, are primarily suitable for parenteral administration, suitable lipophilic solvents or vehicles, such as fatty oils, e.g. sesame oil, or synthetic fatty acid esters, e.g. ethyl oleate or triglycerides, being used, or aqueous injection suspensions which contain viscosity-increasing substances, e.g. sodium carboxymethylcellulose, sorbitol and/or dextran, and if desired also stabilizers.

The dose of the active ingredient can depend on various factors, such as manner of administration, warm-blooded animal species, age and/or individual condition.

Two methods are especially suitable for the measurement of the improvement of the blood circulation in the eye. One is the so-called myograph system for isolated extraocular arteries and the second is the perfusion system in the intact isolated eye, for example a pig's eye.

For the myographic determinations, a short segment (7-8 mm) of the ciliary arteries of fresh eyes, e.g. pig's eyes, is dissected and cut into small rings of thickness 2 mm under a microscope (cf. for this K. Yao et al. *Invest. Ophthalmol. Vis. Sci.* **32**, 1791 - 1798(1991) and M.J. Mulvany, W. Halpern *Circ. res.* **41**, 19 - 26(1977)). During the preparation, the tissue is immersed in modified Krebs-Ringer bicarbonate solution.

These rings (diameter about 200-400 μm) are installed in the myograph system without delay. Such a tissue ring is stretched using two tungsten wires (diameters 30 and 80 μm), the wires being passed through the lumen. One wire is connected to the power transmission, the other to a micromanipulator for adjustment of muscle length. The tissue ring is immersed in a control solution during the experiment (37°; 95 % O_2 ; 5 % CO_2 ; pH = 7.4) and equilibrated for 45 minutes before testing.

The tissue ring is stretched stepwise and at the same time 100 mM KCl is added in each case. The optimum passive tension is then defined as the tension at which the contraction with 100 mM KCl is greatest. For ciliary arteries, an averaged value for the tension of 977 mg \pm 60 mg is found. The active residual tone of this tissue is defined as the difference between the optimum passive tension and the tension after a maximum relaxation with 10^{-6} molar bradykinin. The corresponding tension is 174 mg \pm 38 mg.

In all subsequent experiments, the abovementioned tissue rings are slowly stretched at 100 mg intervals until the optimum is reached.

Before the actual experiment, the endothelial function of such a tissue ring is checked for

- 19 -

intactness. The tissue rings are challenged to contract by using $3 \cdot 10^{-7}$ molar serotonin solution and in the experimental procedure described above. Thereafter it is checked whether a complete relaxation of the contracted tissue ring occurs with $3 \cdot 10^{-7}$ molar bradykinin solution. If this condition is fulfilled, the tissue is held to be intact.

For the perfusion method on the isolated pig's eye, the fatty tissue which surrounds the extraocular muscles is carefully removed. During this preparation, the ocular globe is surrounded with modified Krebs-Ringer bicarbonate solution at 4°C. The ocular globe is then cannulated under the microscope using a polyethylene cannula (600 µm diameter), namely via the common ophthalmic artery.

Following this preparation, the ocular globe is connected via the polyethylene cannula to a Langendorff perfusion system, and in this way supplied at the latest 1 hour after the start of preparation with a filtered and oxygenated Krebs solution which contains 0.5 % albumin.

The catheterization of the common ophthalmic artery guarantees the complete perfusion both of the choroidal and of the retinal tissue and of the small extraocular muscular arteries. The perfused ocular globe is immersed in a thermostated (37°C) control bath with a glass jacket with the cornea pointed downwards. The overflow rate (ml/min) in the bath corresponds to the ophthalmic flow. Before testing a composition, the ocular globe is equilibrated for 20 minutes. In a control experiment, it is shown that the ophthalmic flow in the experimental procedure selected remains constant for about 120 minutes after the start of the perfusion. All tests are therefore carried out within this time frame.

The invention likewise relates to a method of treating normal tension glaucoma, which method comprises administering a therapeutically effective amount of an angiotensin II antagonist or of a pharmaceutically acceptable salt thereof to a patient who needs such a treatment.

The following examples illustrate the invention described above; however, they should not restrict this in its scope in any way. Temperatures are given in degrees Celsius.

Formulation Examples 1, 2 and 3:

A solution, comprising 20 mg of active compound, e.g. valsartan =

(S)-N-(1-carboxy-2-methylprop-1-yl)-N-pentanoyl-N-[2'-(1H-tetrazol-5-yl)biphenyl-4-ylmethyl]amine, can be prepared composed as follows:

- 20 -

Composition:

1)

Active ingredient	20.00 mg
1N NaOH	86.00 mg
Benzalkonium chloride	0.10 mg
Disodium ethylenediaminetetraacetate	0.50 mg
Sorbitol	10.00 mg
Na ₂ HPO ₄ · 2 H ₂ O	9.91 mg
K ₂ HPO ₄	0.44 mg
Water (purity: for inj.) to	1.00 ml

2)

Active ingredient	20.00 mg
1N NaOH	86.00 mg
Macrogol 400	20.00 mg
Benzalkonium chloride	0.10 mg
Disodium ethylenediaminetetraacetate	0.50 mg
Sorbitol	6.00 mg
Na ₂ HPO ₄ · 2 H ₂ O	9.73 mg
K ₂ HPO ₄	0.43 mg
Water (purity: for inj.) to	1.00 ml

3)

Active ingredient	20.00 mg
1N NaOH	86.00 mg
Polyoxyl 35 castor oil	4.00 mg
Benzalkonium chloride	0.10 mg
Disodium ethylenediaminetetraacetate	0.50 mg
Sorbitol	6.00 mg
Na ₂ HPO ₄ · 2 H ₂ O	9.91 mg
K ₂ HPO ₄	0.44 mg
Water (purity: for inj.) to	1.00 ml

For this purpose, the ingredients are added to water and dissolved.

Another A II antagonist or a pharmaceutically acceptable salt thereof can be processed in

an analogous manner, for example as described in the above examples.

Example 4

Determination of the ciliary artery contraction using the myograph method

After it was verified as described above that the endothelial functions of a tissue ring of a ciliary artery are intact, the tissue ring preparation is incubated (contracted) with 10^{-7} M angiotensin II, and then relaxed with various doses of the angiotensin II antagonist valsartan. The results are presented below:

Dose of valsartan	Relative contraction in comparison to 100 mM KCl (in %)
0 (control)	27.9 ± 8.7 (pure angiotensin II effect)
10^{-9} M	18.7 ± 2.2
10^{-8} M	8.1 ± 1.5
10^{-7} M	5.15 ± 1.6
10^{-6} M	0.93 ± 0.4
10^{-5} M	0.9 ± 0.45

The contracting effect of angiotensin II on the ciliary artery is quantitatively abolished using valsartan.

Example 5

A perfused isolated pig's eye is first treated as described above with a control solution and then with a solution comprising angiotensin II (10^{-6} M). After the determination of the angiotensin II effect, valsartan (10^{-5} M) is additionally added on top of the above exposure. The relative ophthalmic flow rate resulting in this way is shown below.


Active ingredient	Decrease in the relative ophthalmic flow (in %)
Control	0.9 ± 2.14
Angiotensin II (10^{-6} M), (AII)	24.5 ± 3.05
AII + valsartan (10^{-5} M)	6.82 ± 3.02

- 22 -


The blood flow-decreasing action of angiotensin II is strongly reduced by valsartan.

WHAT IS CLAIMED IS:


1. The use of an angiotensin II antagonist or of a pharmaceutically acceptable salt thereof in the preparation of a pharmaceutical composition for the treatment of normal tension glaucoma and neurodegenerative processes.
2. The use of a compound according to claim 1, wherein the angiotensin II antagonist is selected from the group consisting of:




(Ia),




(IIa),




(IIIa),




(IVa),




(Va),



(VIa),




(VIb),




(VIIa),


- 25 -




(VIIIa),



(IXa),




(Xa),




(XIa),


- 26 -




(XIIa),



(XIIIa),



(XIVa) and



(XVa).

3. An ophthalmic composition for the treatment of normal tension glaucoma, comprising a therapeutically effective amount of an angiotensin II antagonist or of a pharmaceutically acceptable salt thereof.

4. An ophthalmic composition according to claim 3, wherein the angiotensin II antagonist

- 27 -

is selected from the group of compounds as defined in claim 2.

5. A systemically administrable composition for the treatment of normal tension glaucoma, comprising a therapeutically effective amount of an angiotensin II antagonist or of a pharmaceutically acceptable salt thereof.
6. A systemically administrable composition according to claim 5, wherein the angiotensin II antagonist is selected from the group of compounds as defined in claim 2.
7. A method of treating normal tension glaucoma, which method comprises administering a therapeutically effective amount of an angiotensin II antagonist or of a pharmaceutically acceptable salt thereof to a patient in need of such treatment.
8. A method according to claim 7, wherein the angiotensin II antagonist is selected from the group of compounds as defined in claim 2.

INTERNATIONAL SEARCH REPORT

Intern: Application No
PCT/IB 95/00056

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/00 A61K31/41 A61K31/415 A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,92 10183 (SMITHKLINE BEECHAM PLC) 25 June 1992 see page 20 - page 22 ---	3-6
X	WO,A,93 15732 (CIBA-GEIGY AG) 19 August 1993 see page 2 - page 5; claims 1,4,5 ---	3-6
A	SOC. NEUROSCI. ABSTR., vol. 17, page 810 P. MALLORGA ET AL. 'Angiotensin receptor subtypes in the albino rabbit eye.' abstr 321.12 --- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

7 April 1995

Date of mailing of the international search report

21. 04. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Klaver, T

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IB 95/00056

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	AM. J. HYPERTENS., vol. 4, page 309S R. LOUTZENHISER ET AL. 'Characterization of the renal microvascular effects of angiotensin II antagonist DuP 753.'	
A	GRAEFE'S ARCH. CLIN. EXP. OPHTHALMOL., vol. 230, page 552 R.O.W.BURK ET AL. 'Are large optic nerve heads susceptible to glaucomatous damage at normal intraocular pressure?'	
A	GRAEFE'S ARCH. CLIN. EXP. OPHTHALMOL., vol. 231, page 711 I. GUTMAN ET AL. 'Optic nerve compression by carotid arteries in low-tension glaucoma.'	

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. al Application No
PCT/IB 95/00056

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9210183	25-06-92	AU-A- 9064991	08-07-92
		EP-A- 0561901	29-09-93
		JP-T- 6503335	14-04-94

WO-A-9315732	19-08-93	AU-B- 3722493	03-09-93
		CA-A- 2128324	19-08-93
		EP-A- 0626846	07-12-94
		NO-A- 942756	22-07-94

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.